



ISSN (E): 2320-3862  
ISSN (P): 2394-0530  
NAAS Rating: 3.53  
JMPS 2019; 7(2): 106-116  
© 2019 JMPS  
Received: 11-01-2019  
Accepted: 13-02-2019

#### Tripuresh Dwivedi

a) Patanjali Herbal Research  
Department, Patanjali Research  
Institute, Haridwar,  
Uttarakhand, India

b) Department of Applied  
Medicinal Plants' Sciences, Dev  
Sanskriti Vishwavidyalaya,  
Haridwar, Uttarakhand, India

#### Chandra Kanta

a) Department of Botany, Doon  
(PG.) College of Agriculture,  
Sciences and Technology,  
Dehradun, Uttarakhand, India

b) Department of Biological  
Sciences, College of Basic  
Sciences and Humanities, G B P  
U A & T, Pantnagar, U S Nagar,  
Uttarakhand, India

#### Lalit Raj Singh

Department of Applied  
Medicinal Plants' Sciences, Dev  
Sanskriti Vishwavidyalaya,  
Haridwar, Uttarakhand, India

#### Ishwar Prakash Sharma

a) Patanjali Herbal Research  
Department, Patanjali Research  
Institute, Haridwar,  
Uttarakhand, India

b) Department of Biological  
Sciences, College of Basic  
Sciences and Humanities, G B P  
U A & T, Pantnagar, U S Nagar,  
Uttarakhand, India

#### Correspondence

##### Ishwar Prakash Sharma

a) Patanjali Herbal Research  
Department, Patanjali Research  
Institute, Haridwar,  
Uttarakhand, India

b) Department of Biological  
Sciences, College of Basic  
Sciences and Humanities, G B P  
U A & T, Pantnagar, U S Nagar,  
Uttarakhand, India

## A list of some important medicinal plants with their medicinal uses from Himalayan State Uttarakhand, India

**Tripuresh Dwivedi, Chandra Kanta, Lalit Raj Singh and Ishwar Prakash Sharma**

#### Abstract

Himalayan range has a rich heritage of knowledge on plant based therapy. Medicinal plants play major role in the livelihood from all over the world. Uttarakhand, a Himalayan state of India also depends on the medicinal plants for medicine and traditional therapy. The people from this state use plants for their primary health care system mainly depend on traditional knowledge of medical practices and medicinal herbs. Many of the previous studies on traditional medicine as scientific outputs having traditional claims of effectiveness which are helpful to manage various ailments. In this regards many native medicinal plants listed in article are significantly utilized by the locals, which was need to document. In this scenario we try to document some medicinal plants with their medicinal properties from this state.

**Keywords:** medicinal plants, traditional knowledge, tribal, Uttarakhand

#### Introduction

According to floral statistics of India 2017 hosted by Botanical Survey of India, Kolkata, West Bengal, a total of 2,68,600 flowering plants worldwide among them 18,386 (6.84% of world) existed in India. In India, approximately 3000 plants species are known to have their medicinal properties<sup>[1]</sup>, in another reports a total of 2500 plants are of traditional medicine among them 100 plants used regularly<sup>[2]</sup>. Traditional medical knowledge of plants not only useful for conservation of biodiversity but also useful to healthcare and drug development. Himalaya is a global biodiversity hotspot with much diversified geographical, ecological and evolutionary factors for species diversity which support 18,440 species of plants of which 25.3% is of endemic<sup>[3, 4]</sup>. The Indian state Uttarakhand located in Himalayan hotspot having a huge wild diversity which occupies 17.3% of India's total land area including 92.57% area under hills and 7.43% under plains. Geographically it is located between 28°43'–31°27'N latitudes and 77°34'–81°02'E longitudes. In the border, river Tons separates it from Himachal Pradesh in the north-west, river Kali separates it from Nepal in the east and greater Himalaya is the northern boundary of the state and also the international border with China. About 1748 economically important plants reported from Himalaya<sup>[4]</sup>. The people community mainly Bokshas, Tharus, Bhotias, Van-gujjars, Marchchas, Tolchas, Jaunsaris, Koltas, Gangwal, Banw-rauat, etc. are generally dependent on the wild flora for their own traditional system of therapy<sup>[2, 5]</sup>. Joshi *et al.*<sup>[6]</sup> reported a total of 102 plant species from 48 families are of their ethno-medicinal uses from the four district of the state namely Almora, Champawat, Bageshwar and Pithoragarh. Adhikari *et al.*<sup>[7]</sup> examined the status and distribution pattern of medicinal plants in Wildlife Institute of Dehradun, Uttarakhand and recorded 605 plants from 94 families. This article based on previous studies on medicinal plants of Uttarakhand on this basis authors focused on some important highlights of native medicinal plants which need to conservation and cultivation because these plants are naturally grown in abundance and fast depleting can help the natives to earn their livelihood to some extent. For future perspectives this study, will be helpful to pharmacologist, phytochemist and researcher of this field. Therefore, this study is aimed to manuscript and underlines the importance of traditional knowledge used for the treatment of different diseases in the Himalayan state Uttarakhand, India.

**Table 1:** A list of selected medicinal plants with their ethno-medicinal uses from Uttarakhand, India

Botanical Name (Family)	Local name	Parts used	Ethno-medicinal Uses
<i>Abelmoschus moschatus</i> Medik. Malvaceae	Jangali bhindi	Root, Seed	Fresh roots with young plant of <i>Bombax ceiba</i> used in the treatment of leucorrhoea in female and sexual impotency in male <sup>[2]</sup> . Seeds are diuretic, aphrodisiac, ophthalmic, cardiotoxic, digestive, stomachic, constipating, carminative, stimulant, antispasmodic, cardiac debility, cough, asthma, bronchitis, hyperdipsia, burning sensation, nausea, dyspepsia, flatulent colic, diarrhea, strangury, gonorrhoea, spermatorrhea, calculi, halitosis, ptyalism, vomiting and other neural disorders, leukoderma, and general debility <sup>[8, 9]</sup> .
<i>Acorus calamus</i> L. (Acoraceae)	Baj	Root	Headache <sup>[10]</sup> ; Anti-bacterial, anti-helminthic <sup>[11]</sup> ; Fever, asthma, bronchitis, cough, digestive problems (gas, bloating, colic) <sup>[12]</sup> .
<i>Adenostemma lavenia</i> (L.) Kuntze Asteraceae	Jangli-jira	Flower, Leaf	Flowers used for increasing saliva so useful in treatment of mouth dryness <sup>[2]</sup> . Leaves paste use in wound healing <sup>[13]</sup> .
<i>Aegle marmelos</i> (L.) Corrêa (Rutaceae)	Bel	Fruit, Root	Astringent, antidiarrheal, antidysenteric, demulcent, antipyretic, laxative <sup>[10, 14, 15]</sup> .
<i>Aesculus indica</i> (Wall. ex Cambess.) Hook. (Sapindaceae)	Panker	Fruit	Rheumatism (warm paste applies on affected part) <sup>[10, 16]</sup> .
<i>Ageratum conyzoides</i> L. Asteraceae	Goatweed	Leaf	Leaves paste is very useful in wound healing <sup>[2]</sup> , treatment of leprosy, diarrhea, dysentery, intestinal colic, rheumatism, fever <sup>[17]</sup> .
<i>Allium cepa</i> L. (Amaryllidaceae)	Pyaj	Bulb	Roasted warm bulb is eaten in abdominal pain and skin diseases <sup>[10]</sup> and also used in malaria, diarrhea, wounds with antimicrobial, antioxidant properties <sup>[18]</sup> .
<i>Amelanchier canadensis</i> (L.) Medik. Rosaceae	Indian wild pear	Fruit	The juice of the ripe fruit is used in the treatment of diarrhea <sup>[19, 20]</sup> .
<i>Anaphalis triplinervis</i> Sims ex C.B. Clarke (Asteraceae)	Bakal	Leaf	Used in wounds and skin diseases <sup>[10]</sup> important in flu, fever, nausea and eye infections <sup>[21]</sup> .
<i>Anisomeles indica</i> (L.) Kuntze Lamiaceae	Catmint	Leaf	Fresh leaves as well as greenish parts contain volatile oil used in treatment of cough and cold <sup>[2]</sup> , chronic rheumatism, psoriasis, snake bites <sup>[22]</sup> .
<i>Asparagus curillus</i> Buch.-Ham. ex Roxb. (Asparagaceae)	Safed Musli	Root	Root powder is administered as sexual tonic <sup>[10]</sup> along with dysuria, diabetes and dysentery <sup>[23]</sup> .
<i>Asparagus racemosus</i> Willd. (Asparagaceae)	Satawari kairua, Katerwali vel	Root, Rhizome	Root powder administered orally as aphrodisiac <sup>[10]</sup> ; also used in stomachache, urinary disorders, Nervous disorders, inflammation, liver diseases, infectious diseases <sup>[24, 25]</sup> , ulcers, cancer <sup>[26]</sup> .
<i>Berberis aristata</i> DC. (Berberidaceae)	Kambal	Root	Root extract used in eye diseases, diarrhea, jaundice, skin diseases, syphilis, chronic rheumatism, urinary disorders <sup>[10, 27]</sup> , diabetes, jaundice and Gastro intestinal problem <sup>[28]</sup> .
<i>Berberis chitria</i> Buch.-Ham. ex Lindl. (Berberidaceae)	Kilmorha	Root	Root extract is very useful in higher fever, jaundice, diabetes, stomach disorders, rheumatism, eye, ear and skin disease <sup>[10, 29]</sup> .
<i>Bergenia ciliata</i> (Haw.) Sternb. (Saxifragaceae)	Sirparha	Root	Root powder is very useful in kidney stones and ulcers <sup>[10, 28, 30]</sup> .
<i>Boenninghausenia albiflora</i> (Hook.) Rchb. ex Meisn. (Rutaceae)	Pissumar	Whole plant	External application of whole plant juice cure headache and eyes pain <sup>[10]</sup> ; also shows various effects as hepatoprotective, antioxidative, anti-inflammatory and immunomodulating <sup>[31]</sup> .
<i>Boerhavia diffusa</i> L. (Nyctaginaceae)	Punarnava	Root	Juice of fresh roots is used as eye drops <sup>[10]</sup> ; very useful tonic to liver problem, jaundice, asthma with antimicrobial activity <sup>[32, 33]</sup> .
<i>Bombax ceiba</i> L. (Malvaceae)	Saimul	Leaf, Root	Leaves paste applies over wounds with good property of pain killer and improve gynecological disorders <sup>[10, 34]</sup> . The formulation of root with leaves of <i>Dalbergia sissoo</i> is used in sexual impotency <sup>[2]</sup> .
<i>Boswellia serrata</i> Roxb. (Burseraceae)	Sallai	Bark	Hot decoction of bark used in wound <sup>[10]</sup> , chronic inflammatory diseases <sup>[35]</sup> ; diarrhea <sup>[36]</sup> .
<i>Buchanania cochinchinensis</i> (Lour.) M.R. Almeida (Anacardiaceae)	Achar	Leaf	Very effective in dental pain when gargle the juice twice daily <sup>[10]</sup> ; asthma <sup>[37]</sup> ; cancer <sup>[38]</sup> .
<i>Butea monosperma</i> (Lam.) Kuntze Fabaceae	Dhak, Palash	Flower, Gum	The flower extract with water useful in gastritis <sup>[2]</sup> . Gum from bark known as Bengal Kino useful in treatment of ulcer, diarrhea, antioxidative, anti-inflammatory, hepatoprotective and anti-cancer activities <sup>[39]</sup> .
<i>Byttneria herbacea</i> Roxb. (Malvaceae)	Kamraj	Root	Root paste used in fever and leucorrhoea <sup>[10]</sup> .
<i>Cajanus scarabaeoides</i> (L.) Thouars (Fabaceae)	Chowkhara	Leaf	Crushed or rubbed leaves applied in throat disease, chewed in tonsillitis <sup>[10]</sup> ; Hepatoprotective and immense medicinal potential <sup>[40, 41]</sup> .
<i>Callicarpa macrophylla</i> Vahl. (Lamiaceae)	Daya	Fruit	Very helpful in urinary disorders, skin problem, diarrhea, defecation and hair care <sup>[10, 42, 43]</sup> .
<i>Calotropis gigantea</i> (L.) W.T. Aiton (Apocynaceae)	Aak	Leaf	Warmed fresh leaves used in wound, toothache, earache, epilepsy, diarrhea and mental disorders <sup>[10, 44]</sup> .

<i>Carissa spinarum</i> L. (Apocynaceae)	Karunda	Root	Used in rheumatism, having strong purgative property so used as one of the ingredients in some purgative preparations <sup>[20, 45]</sup> .
<i>Celastrus paniculatus</i> Willd. (Celastraceae)	Malkangani	Fruit, Seed	Boiled solution of fruits and seeds with oil use in piles, gout, rheumatism, cold, dysentery, diarrhoea, leprosy, snake bite, wound, anti-spasmodic, carminative and antihelmintic properties and also used for treatment of epilepsy, mental ailments, chronic diarrhea, dysentery, bronchial catarrh, intermittent fevers and tumors <sup>[46]</sup> ; pneumonia <sup>[47]</sup> ; leucorrhoea <sup>[48]</sup> .
<i>Cleome viscosa</i> L. (Cleomaceae)	Kumkum	Leaf	Leave's juice very useful to ear diseases <sup>[10]</sup> ; anthelmintic, antiseptic, carminative, antiscorbutic, sudorific, febrifuge, and cardiac problems <sup>[49]</sup> .
<i>Commelina benghalensis</i> L. Commelinaceae	Kanchara	Leaf	Paste of leaves used for curing swelling as honeybee biting <sup>[2]</sup> .
<i>Commelina paludosa</i> Blume Commelinaceae	Kankowa	Whole plant	Decoction of plants with water used in treatment of sexual impotency <sup>[2]</sup> .
<i>Cordia myxa</i> L. Boraginaceae	Lasura	Fruit	Fruit are very useful in common cold and cough <sup>[2]</sup> .
<i>Corylus colurna</i> L. (Betulaceae)	Bhotia badam	Nut	Direct use of nut or its tonic use as diuretic, aphrodisiac <sup>[50]</sup> .
<i>Cydonia oblonga</i> Mill. (Rosaceae)	Bihi	Leaf, Seed	Leaves having phenolic compounds as antioxidants properties, Vit-E, carotenoids, L-ascorbic acid and other organic acids <sup>[51, 52]</sup> ; in folk medicine for their sedative, antipyretic, anti-diarrheic and antitussive properties along with treatment of various skin diseases <sup>[53, 54]</sup> . Seeds used in diarrhea, dysentery, cough, sore throat and bronchitis <sup>[55]</sup> , intestinal colic and constipation <sup>[56]</sup> and also used for allergic rhinitis and asthma <sup>[20, 57]</sup> .
<i>Datura metel</i> L. (Solanaceae)	Dhatur	Leaf	Boil and warm leaves used in affected part for antibiotics, antibacterial, antimicrobial activity <sup>[10, 58]</sup> .
<i>Datura stramonium</i> L. (Solanaceae)	Dhatura	Flower	Flower's juice used in earache <sup>[10]</sup> having analgesic and antiasthmatic activities <sup>[59]</sup> .
<i>Dicliptera bupleuroides</i> Nees (Acanthaceae)	Soriul	Seed, Leaf	Decoction of seeds and leaves with water very useful in dysentery <sup>[10]</sup> ; respiratory and urinary tract infections, digestive disorders, sexual problems, skin diseases, liver ailments, wound healing, blood vomiting, malaria, piles, epilepsy, jaundice, gummosis, round worms, hepatitis <sup>[60]</sup> .
<i>Drimia indica</i> (Roxb.) Jessop (Asparagaceae)	Banpyaja	Bulb	Juice of bulb is used in cough, bronchitis, nematode infection, pyrexia <sup>[10]</sup> dropsy, respiratory ailment, bone and joint complications, skin disorders, epilepsy and cancer <sup>[61]</sup> .
<i>Euonymus tingens</i> Wall. (Celastraceae)	Kusum	Whole plant	The powder mixture of 10gm with opium seeds (10gm), almond (5gm), kali mirch (5gm) and bansh mishri (10gm) used orally in kidney stone and eye diseases <sup>[10, 62]</sup> .
<i>Ficus auriculata</i> Lour. (Moraceae)	Timil	Whole plant	Gastrointestinal problems treated by using 50-100ml fresh leaves juice when used regularly upto 10 days <sup>[63]</sup> . Bark and root show hypoglycemic and anthelmintic activity <sup>[64, 65]</sup> . Fruit extracts exhibits anti-tumour activity <sup>[20, 66]</sup> .
<i>Ficus auriculata</i> Lour. Moraceae	Timul, Timil	Whole plant	Gastrointestinal problems treated by using 50-100 ml fresh juice of leaves with water for about 10 days <sup>[63]</sup> . Bark and root show hypoglycaemic and anthelmintic activity <sup>[64, 65]</sup> . The extract inhibits insulinase activity from liver and kidney. Fruit extracts exhibits anti-tumour activity <sup>[66]</sup> . Leaves exhibit hypotensive activity <sup>[20, 67]</sup> .
<i>Ficus benghalensis</i> L. Moraceae	Bargad	Fruit, Bark	The milky latex of fruit with sugar used in treatment of sexual impotency <sup>[2]</sup> . Leaf, fruit, bark are useful in central nervous system, endocrine system, gastrointestinal tract, reproductive system, respiratory system and infectious disorders <sup>[68]</sup> .
<i>Ficus palmata</i> Forssk. (Moraceae)	Bedu	Fruit	Fruits are beneficial in lung and bladder diseases. Along with these are good source of minerals, phosphorus and a small amount of Vit. C <sup>[69]</sup> ; the latex is useful in wound healing <sup>[28]</sup> .
<i>Geranium wallichianum</i> D. Don ex Sweet (Geraniaceae)	Laljari	Root	Root extract useful in backache, gout, bone strengthening, hepatitis, liver problems, premature delivery <sup>[30, 70]</sup> .
<i>Glycosmis pentaphylla</i> (Retz.) DC. Rutaceae	Ban Nimbu	Seed	Seeds of plant used in curing of vomiting <sup>[2]</sup> .
<i>Grevillea robusta</i> A. Cunn. ex R.Br. Proteaceae	Silver Oak	Leaf	Paste of fresh leaves with vegetable ghee and coconut oil very useful in burning and burning sensation <sup>[2]</sup> .
<i>Gymnema sylvestre</i> (Retz.) R.Br. ex Sm. (Apocynaceae)	Aphe	Leaf	Boiled leaves with til-oil when applied externally 3-4 drops twice per day used for eyes to reduce pain. Direct use of milky juice from crushed fresh leaves when applied once daily for 3 day also helpful in eye problems <sup>[10]</sup> along with antidiabetic properties <sup>[71]</sup> .
<i>Gymnosporia spinosa</i> (Blanco) Merr. & Rolfe (Celastraceae)	Vaikal	Leaf	Paste of leaves useful in Bala-shosha (a condition in which child is emaciated) when applied and rubbed on back <sup>[10]</sup> .
<i>Hedychium spicatum</i> Sm. (Zingiberaceae)	Kapurkachari	Root	Root powder is very useful in neuro muscular disorders and body pain along with anticancerous and antimicrobial activities <sup>[10, 72]</sup> .
<i>Helicteres isora</i> L. (Malvaceae)	Atanda	Root	Orally use of crushed roots used in excessive appetite, empena and stomachic <sup>[10, 73]</sup> ; having antioxidant and anticancer properties <sup>[74]</sup> .
<i>Holarrhena pubescens</i> Wall. ex G. Don	Dhodhi	Latex	Useful in ring worm when latex applied on affected part <sup>[10]</sup> also useful in constipation, colic, and diarrhea <sup>[75]</sup> .

(Apocynaceae)			
<i>Hydrocotyle sibthorpioides</i> Lam. (Araliaceae)	Brahmni	Leaf	Leaf juice is administered orally in Manasdaurbalya (mental disorder) <sup>[10]</sup> ; also used in various problems as dysentery, diarrhoea, piles, rheumatism, digestive, diuretic, vermifuge, menstrual problem <sup>[76]</sup> .
<i>Hygrophila auriculata</i> (Schumach.) Heine (Acanthaceae)	Talmakhana	Seed	Root is used for rheumatism, inflammation, jaundice, hepatic obstruction, pain <sup>[77]</sup> ; whole plant extract used in treatment of urinary infection, gout, hepatic obstruction and as a diuretic <sup>[78]</sup> .
<i>Hypericum japonicum</i> Thunb. (Hypericaceae)	Chhingewali	Whole plant	Plant powder is used as snuff for nose disease <sup>[10]</sup> ; along with hepatoprotective activity, protecting liver injury and hepatotoxicity <sup>[79]</sup> .
<i>Jacobaea nudicaulis</i> (Buch.-Ham. ex D.Don) B.Nord. (Asteraceae)	Nil kanthi, Ratpatia	Whole plant	Plant juice applied in eye diseases, wound, fever and some skin diseases <sup>[10, 62]</sup> .
<i>Juniperus communis</i> L. (Cupressaceae)	Hapusha	Green wood	Excessive appetite and smoke of green wood is used in tuberculosis and respiratory disease <sup>[10, 80]</sup> ; along with antioxidant and antimicrobial activities <sup>[81]</sup> .
<i>Justicia adhatoda</i> L. (Acanthaceae)	Bhasma	Leaf, Flower	Juice of leaves and flowers used in cough, fever, coryza, respiratory problems <sup>[10, 82, 83]</sup> .
<i>Kalanchoe pinnata</i> (Lam.) Pers. (Crassulaceae)	Sanjwanboata	Leaf	Paste of leaves is used over wounds for healing <sup>[10]</sup> .
<i>Lablab purpureus</i> (L.) Sweet (Fabaceae)	Chimi	Whole plant	Fresh juice is dropped in earache <sup>[10]</sup> and used as antidiabetic, antiinflammatory, analgesic, antioxidant, cytotoxic, hypolipidemic, antimicrobial, insecticidal, hepatoprotective, antilithiatic, antispasmodic <sup>[84]</sup> .
<i>Lansea coromandelica</i> (Houtt.) Merr. (Anacardiaceae)	Gunjhanganj	Bark	Paste of bark is applied on cuts to stop bleeding and wound healing <sup>[10]</sup> ; while juice use in jaundice and liver pain <sup>[85, 86]</sup> .
<i>Leucas cephalotes</i> (Roth) Spreng. (Lamiaceae)	Guma	Whole plant	Orally use of whole plant used in fever and decoction <sup>[10]</sup> ; Asthma, cough <sup>[87]</sup> .
<i>Lilium polyphyllum</i> D.Don (Liliaceae)	Kakoli	Root	Tonic and decoction of root used as refrigerant, galactagogue, expectorant, aphrodisiac, diuretic, antipyretic and tonic <sup>[8, 88]</sup> .
<i>Litsea glutinosa</i> (Lour.) C.B. Rob. Lauraceae	Maida	Bark	The bark is used to relieve pain, arouse sexual power, produce a soothing effect on the body and arrest bleeding. Bark paste is applied to bind fractured limbs <sup>[2, 89]</sup> .
<i>Melia azedarach</i> L. (Meliaceae)	Dekrain	Root	Paste of root is applied in headache <sup>[10]</sup> ; it has also various activities as anthelmintic, antilithic diuretic, astringent, stomachic, anticancer, antimalarial, analgesic, anti-inflammatory activities <sup>[90]</sup> .
<i>Mentha longifolia</i> (L.) L. (Lamiaceae)	Piper mint	Leaf	Juice or powder of leaves uses in common fever, coryza, rheumatism, dysentery, Dyspepsia <sup>[10, 91]</sup> .
<i>Micromeria biflora</i> (Buch.-Ham. ex D.Don) Benth. (Lamiaceae)	Masipatha	Whole plant	Whole plant is administered orally with milk to control fever, Sciatica, arthritis, skin disease <sup>[10, 92]</sup> .
<i>Mimosa rubicaulis</i> Lam. (Fabaceae)	Shikanta	Leaf	Leaf paste used in skin disease when applied directly on affected part <sup>[10]</sup> ; along with various other problems as bronchitis, cholera, cough, dyspepsia, fever, jaundice, smallpox, syphilis and tuberculosis <sup>[55]</sup> .
<i>Morella esculenta</i> (Buch.-Ham. ex D.Don) I.M.Turner (Myricaceae)	Kafal, Kaphal	Fruit, Bark	Specified decoctions of fruit, stone and bark are claimed to be beneficial in cardiac debility, edema and hemoptysis; wax from fruit covering used for ulcer healing <sup>[93-96]</sup> ; along with anti-inflammatory activity and anti tumour activities <sup>[97]</sup> .
<i>Morina longifolia</i> Wall. ex DC. (Caprifoliaceae)	Vishkanya	Whole plant	Oral administration of plant use for wound healing along with various activities as antimicrobial, cytotoxic, antiulcer, hypoglycemic and hypotensive <sup>[10, 98]</sup> .
<i>Nyctanthes arbor-tristis</i> L. (Oleaceae)	Siyari	Leaf	Juice of leaves is applied on affected part to cure skin diseases <sup>[10]</sup> ; along with immunotoxic, antiallergic, antihistaminic, purgative, antibacterial and cytotoxicity, antipyretic and ulerogenic, anti -Inflammatory activity <sup>[99]</sup> .
<i>Origanum vulgare</i> L. (Lamiaceae)	Vantulsi	Whole plant	Urinary disorder can be cure by oral decoction of whole plant <sup>[10]</sup> ; various activities as antifungal <sup>[100]</sup> and anti-hyperglycemic activity <sup>[101]</sup> also reported.
<i>Oxalis corniculata</i> L. (Oxalidaceae)	Chilmora, Salmosi	Whole plant	Juice of plant is used to cure various eye diseases such as motiabinda <sup>[10]</sup> ; along with it used in bacterial diseases, dysentery, diarrhea, skin disease <sup>[102]</sup> .
<i>Phyllanthus emblica</i> L. (Phyllanthaceae)	Amla	Whole plant	Fruit are rich in polyphenols, minerals and regarded as one of the richest source of Vit. C <sup>[103]</sup> . Therapeutically it has energy refilling potential, aperient, antibacterial, antifungal, antiviral activities <sup>[104]</sup> , along with gonorrhoea, analgesic and skin fairness <sup>[93]</sup> and to stop nausea and vomiting, antitumour and hepatoprotective activity <sup>[20, 104]</sup> .
<i>Piper longum</i> L. (Piperaceae)	Pippali	Fruit	Powder of fruits is administered orally to cure cough, respiratory tract bronchitis, asthma, analgesic, muscular pains, inflammation, hematinic, carminative <sup>[10, 105]</sup> .
<i>Plantago major</i> L. (Plantaginaceae)	Isabgol	Seed	Seed powder is useful in dysentery, wound healing, anti-inflammation, analgesic, antioxidant, antiulcer genic activity <sup>[10, 106]</sup> .
<i>Pleurolobus gangeticus</i> (L.) J.St.-Hil. (Fabaceae)	Salparni	Leaf	Paste of fresh leaves is applied on affected part for wound healing and used for toothache, chest pains, fungal infections <sup>[10, 107]</sup> .
<i>Plumbago zeylanica</i> L. (Plumbaginaceae)	Chitavar	Root, Fruit	Paste of root and fruit is applied on affected part for wound healing <sup>[10]</sup> and also having anti-atherogenic, cardiogenic, hepatoprotective and neuroprotective

			properties <sup>[108]</sup> ; very useful to rheumatic pain, dysmenorrhea, carbuncles, contusion of the extremities, ulcers and elimination of intestinal parasites <sup>[109]</sup> .
<i>Polygonatum cirrhifolium</i> (Wall.) Royle (Asparagaceae)	Mahameda	Root	Tonic and root powder is administered orally as anti-inflammatory, analgesic, antidiarrheal, antimicrobial, antioxidant and antimalarial <sup>[10, 110]</sup> .
<i>Polygonatum verticillatum</i> (L.) All. (Asparagaceae)	Meda	Rhizome	Tonic and powder of rhizomes is administered orally as aphrodisiac, emollient, cardio tonic, carminative, sialagogue, stimulant <sup>[10, 111]</sup> .
<i>Prunus armeniaca</i> L. (Rosaceae)	Khubani	Fruit	The fruit having high in carotene and Vit. C, provides a valuable source of food. The amygdalin a chemical extracted from apricot kernels used as an alternative treatment for cancer <sup>[112]</sup> along with having antioxidant and antimicrobial properties <sup>[113]</sup> .
<i>Punica granatum</i> L. (Lythraceae)	Dadim	Fruit	Fruit is very useful to worm infestation with antimicrobial, antioxidant, anticancer, anti-inflammatory and anti-cancerous activities <sup>[10, 114]</sup> .
<i>Pyracantha crenulata</i> (D.Don) M.Roem. (Rosaceae)	Ghigharu	Fruit, Bark	Fruits used as preservative. Medicinally it has cardio-tonic, coronary vasodilator and hypertensive properties and used in cardiac failure, myocardial weakness, paroxysmal tachycardia, hypertension, arteriosclerosis and Burgor's disease. Rejuvenation property, reduce joint pains. The bark used in heavy bleeding during menstrual cycles, malarial fever <sup>[20, 115, 116]</sup> .
<i>Ribes nigrum</i> L. (Grossulariaceae)	Blackcurrant	Leaf	The extracted oil and juice useful as an antioxidant source and in treating rheumatoid arthritis and night and fatigue-related visual impairment, antimicrobial and anticancer <sup>[20, 117, 118]</sup> .
<i>Ricinus communis</i> L. (Euphorbiaceae)	Ein	Leaf	Warm fresh leaves are smeared and applied on affected part for wound healing with antimicrobial, toxicological and anti-inflammatory activity <sup>[10, 119]</sup> .
<i>Rosa canina</i> L. (Rosaceae)	Dog Rose	Leaf, Fruit	The plant has high antioxidants and vit-C level, used to make syrup, tea and marmalade. The fruits have been used internally as tea for treatment of viral infections and disorders of the kidneys and urinary tract <sup>[20, 120, 121]</sup> .
<i>Rosa moschata</i> Herrm. (Rosaceae)	Kunja	Leaf	Leaves juice is used as nasal drops to control hemorrhage <sup>[10]</sup> ; useful to control stomach disorder <sup>[122, 123]</sup> .
<i>Roylea cinerea</i> (D.Don) Baill. (Fabaceae)	Kaural	Leaf	Fresh leaves juice is useful in diabetes, mouth diseases and throat diseases <sup>[10]</sup> ; including antioxidant and antimicrobial activities <sup>[124]</sup> to provide strength to liver and protect skin from infection <sup>[125]</sup> .
<i>Rubus ellipticus</i> Sm. (Rosaceae)	<b>Lalanchu</b>	Fruit	It has good antioxidant properties to providing free energy for the people who are travelling mountains <sup>[19, 126]</sup> .
<i>Rumex hastatus</i> D. Don (Polygonaceae)	Amloraha	Root, Leaf	Uses in decoction when administered orally for skin diseases, leaf juice is good for abdominal colic <sup>[10]</sup> including Antioxidant and anticholinesterase <sup>[127]</sup> and anti-tuberculosis properties <sup>[128]</sup> .
<i>Rumex nepalensis</i> Spreng. (Polygonaceae)	Jangali palak	Leaf	Juice or powder of leaves is useful in abdominal colic and skin diseases <sup>[10]</sup> with wound healing and anti-allergic properties <sup>[28]</sup> .
<i>Rumex vesicarius</i> L. (Polygonaceae)	Chalmora	Leaf	Leaves are good source of antioxidants with good antibacterial properties <sup>[129, 130]</sup> .
<i>Satyrium nepalense</i> D.Don (Orchidaceae)	Mishri	Root	Energetic tonic from roots cure various fever including antibacterial, anti-pneumonia properties <sup>[131]</sup> .
<i>Sida cordifolia</i> L. (Malvaceae)	Denusha	Root, Bark	Tonic of stem bark and root powder is given in general debility as stomatitis, blenorhea, asthmatic bronchitis, nasal congestion <sup>[132]</sup> .
<i>Smilax aspera</i> L. (Smilacaceae)	Kukundara	Root	Paste of root is given orally with water to control diarrhea <sup>[10, 133]</sup> .
<i>Solanum lasiocarpum</i> Dunal (Solanaceae)	Chitrika	Leaf, Root	Seven leaves are to be placed on the head during sleeping and this process is to be continued for three nights to control fever and decoction of root is administered orally <sup>[10]</sup> .
<i>Solanum nigrum</i> L. Solanaceae	Black Nightshade	Whole plant	It has expectorant, analgesic, sedative, diaphoretic properties. Its external application cures skin diseases and gives relief in burns, itching, pain etc. Leaves juice used in earache <sup>[20, 134-138]</sup> .
<i>Solanum violaceum</i> Ortega (Solanaceae)	Brahati	Fruit	Four ripe fruits are taken orally at a time to control cough <sup>[10]</sup> along with various properties such as hypertension, poisonous, insect bites <sup>[139]</sup> .
<i>Solena heterophylla</i> Lour. (Cucurbitaceae)	Gulakhari	Tuber	Tubers are eaten as vegetables to control hiccough, asthma, nausea and incisions; useful to control malaria, diabetes, toothache and various disorders as gastrointestinal, respiratory and vascular disorders <sup>[140]</sup> .
<i>Swertia chirayita</i> (Roxb.) H.Karst. (Gentianaceae)	Chirata	Whole plant	Decoction of whole plant is administered orally to control fever, skin disease, dyspepsia, diarrhea with antipyretic, antifungal, hypoglycemic properties <sup>[44, 141, 168]</sup> .
<i>Syzygium cumini</i> (L.) Skeels Myrtaceae	Jamun	Bark	Bark extract useful diabetes <sup>[2]</sup> , sore throat, bronchitis, asthma, thirst, biliousness, dysentery and ulcers <sup>[142]</sup> .
<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn. Combretaceae	Arjun	Bark	Bark is very useful to treatment of pneumonia <sup>[2]</sup> , fractures, ulcers, hepatic and shows hypocholesterolemic, antibacterial, antimicrobial, antitumoral, antioxidant, antiallergic and antifeedant, antifertility and anti-HIV activities <sup>[143, 144]</sup> .
<i>Thymus serpyllum</i> L. (Lamiaceae)	Van Ajwain	Leaf	Leaves used as spice to control dyspepsia <sup>[10]</sup> .
<i>Trapa natans</i> var. <i>bispinosa</i> (Roxb.) Makino (Lythraceae)	Singhara	Nut	Nuts are very useful in diarrhea, dysentery, cardiac diseases, blood pressure with having antimicrobial, cytotoxic activities <sup>[145]</sup> .

<i>Urena lobata</i> L. (Malvaceae)	Bara Mamas	Whole plant	Tonic and paste of whole plant is administered orally with milk to control urinary problems and sexual transmitted diseases [146].
<i>Urtica ardens</i> Link (Urticaceae)	Shishuna	Leaf	Leaves are useful in bone fracturing, cough, cold, digestive problems, fever, headache, skin infection [147].
<i>Vachellia nilotica</i> (L.) P.J.H.Hurter & Mabb. Fabaceae	Babul	Whole plant	Leaves have chemoprventive, anitmutagenic, anti-bacterial, anticancer, astringent, anti-microbial, anti-ulcers, anti-inflammatory activities [148, 149]. The roots are used in cancers, tuberculosis and indurations of liver and spleen [48]. Bark is very useful as antibacterial, antioxidant, anti-mutagenic, cytotoxic, astringent, acrid cooling, styptic, emollient, anthelmintic, aphrodisiac, diuretic, expectorant, emetic, nutritive, in hemorrhage, wound ulcers, leprosy, leucoderma, small pox, skin diseases, biliousness, burning sensation, toothache, leucoderma, dysentery and seminal weakness [48,150-155].
<i>Verbascum thapsus</i> L. (Scrophulariaceae)	Kalber	Leaf	Boiled leaves are applied over inflamed parts to cure headache, inflammatory diseases [10]; also used to control asthma, spasmodic coughs, diarrhea, Pulmonary problems [156].
<i>Vigna mungo</i> (L.) Hepper (Fabaceae)	Urad dal	Fruit	Paste of fruits applied over fractured bone to join [10]; also helpful to control urinary tract infections, sexually transmitted diseases [85, 157].
<i>Vitex negundo</i> L. (Lamiaceae)	Nirgundi	Leaf	A syrup of leaves with rice and water given orally to treat gonorrhoea [10] along with anti-inflammatory, anticonvulsant, antioxidant, bronchial relaxant, hepatoprotective properties [158].
<i>Vitis vulpina</i> L. (Vitaceae)	Jungle Angoor	Leaf Fruit	Unripe grapes used for treating sore throats; raisins useful to control tuberculosis while ripe grapes used for the treatment of cancer, cholera, smallpox, nausea, skin & eye infections, kidney & liver diseases [20, 159-161].
<i>Zanthoxylum armatum</i> DC. (Rutaceae)	Timur	Leaf	Regular use of leaves' Pakora up to 3-4 days very useful to control allergy with carminative, stomachic, anthelmintic activities [10, 162].
<i>Ziziphus jujuba</i> Mill. (Ramanaceae)	Ber	Whole plant	Delicious fruits used to increase weight, stamina, improves muscular and liver strength along with diuretic, emollient and expectorant properties. Dried fruits useful as anticancer, pectoral, refrigerant, sedative, stomachache, blood purifier. Seeds are used internally in the treatment of palpitations, insomnia, nervous exhaustion, night sweats and excessive perspiration A decoction of the root has been used in the treatment of fever, wound and ulcer treatment. The leaves are applied in liver troubles, asthma and fever [20, 163].

### Medicinal plants with their herbal uses

A total of 111 medicinal plants from the Himalayan Indian state of Uttarakhand have been listed in table with their medicinal properties. All botanical information of these plants evaluated and confirm from the Plants of the World Online database site.

### Conclusion

Medicinal plants are the natural health care to the people. Their primary cure of diseases is based upon deep observation of nature and their understanding of traditional knowledge of medical practices. Local people of Uttarakhand heavily use these traditionally easily available medicinal plants for health which are less expensive without side effects. The plants now a days going to extinct due to development activities, population explosion, impact of tourism, deforestation, etc. which need to conserve for biodiversity, natural and local aspect.

### References

1. Prakasha HM, Krishnappa M, Krishnamurthy YL, Poornima SV. Folk medicine of NR PuraTaluk in Chikamagalur district of Karnataka. Indian Journal of Traditional Knowledge. 2010; 9(1):55-60.
2. Joshi B, Pant SC. Ethnobotanical study of some common plants used among the tribal communities of Kashipur, Uttarakhand. Indian Journal of Natural Products and Resources. 2012; 3(2):262-266.
3. Singh DK, Hajra PK. Floristic diversity. In: Gujral GS, Sharma V (Eds), Changing Perspective of Biodiversity Status in the Himalaya. British Council Division, British High Commission Publ Wildlife Youth Services, New Delhi, 1996.
4. Samant SS, Dhar U, Palni LMS. Medicinal Plants of Indian Himalaya: Diversity Distribution Potential Value.

Almora: G.B Pant Institute of Himalayan Environment and Development, 1998.

5. Gaur RD. Traditional dye yielding plants of Uttarakhand, India. Natural Product Radiance. 2008; 7(2):154-165.
6. Joshi Y, Joshi AK, Prasad N, Juyal D. A review on *Ficus palmate* (Wild Himalayan Fig). Journal of Psychopharmacology. 2014; 3(5):374-377.
7. Adhikari BS, Babu MM, Saklani PL, Rawat GS. Medicinal Plants Diversity and their Conservation Status in Wildlife Institute of India (WII) Campus. Ethnobotanical Leaflets. 2010; 14(1):46-83.
8. Harborne JB. Indian Medicinal Plants, A compendium of 500 species Journal of Pharmacy and Pharmacology. 1994; 46(11):635.
9. Pawar AT, Vyawahare NS. Protective effect of ethyl acetate fraction of *Biophytum sensitivum* extract against sodium oxalate-induced urolithiasis in rats. Journal of Traditional and Complementary Medicine. 2017; 7(4):476-486.
10. CCRS. An appraisal of Tribal- folk medicines. Vijay nagar, New Delhi, 1999.
11. McGaw LJ, Jäger AK, van Staden J, Houghton PJ. Antibacterial effects of fatty acids and related compounds from plants. South African Journal of Botany. 2002; 68(4):417-423.
12. Balakumbahan R, Rajamani K, Kumanan K. *Acorus calamus*: An overview Journal of Medicinal Plants Research. 2010; 4(25):2740-2745.
13. Prasad AGD, Shyma TB, Raghavendra MP. Plants used by the tribes for the treatment of digestive system disorders in Wayanad district, Kerala. Journal of Applied Pharmaceutical Science. 2013; 3(8):171-175.
14. Kesari AN, Gupta RK, Singh SK, Diwakar S, Wata G. Hypoglycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats.

- Journal of Ethnopharmacology. 2006; 107(3):374-379.
15. Kala CP. Ethnobotany and ethno conservation of *Aegle marmelos* (L), Correa. Indian Journal of Traditional Knowledge. 2006; 5(4):537-540.
  16. Rajasekaran A, Singh J. Ethnobotany of Indian horse chestnut (*Aesculus indica*) in Mandi district, Himachal Pradesh. Indian Journal of Traditional Knowledge. 2009; 8(2):285-286.
  17. Kamboj A, Saluja AK. *Ageratum conyzoides* L.: A review on its phytochemical and pharmacological profile. International Journal of Green Pharmacy. 2008; 2:59-68.
  18. Okusa PN, Penge O, Devleeschouwer M, Duez P. Direct and indirect antimicrobial effects and antioxidant activity of *Cordia gillettii* De Wild (Boraginaceae). Journal of Ethnopharmacology. 2007; 112(3):476-481.
  19. Manandhar NP. Plants and People of Nepal. Timber Press Oregon. 2002; 7(12):599.
  20. Sharma IP, Kanta C, Semwal SC, Goswami N. Wild Fruits of Uttarakhand (India): Ethnobotanical and Medicinal Uses. International Journal of Complementary & Alternative Medicine. 2017; 8(3):1-8.
  21. Khan B, Abdulkadir A, Qureshi R, Mustafa G. Medicinal uses of plants by the inhabitants of Khunjerab National Park, Gilgit, Pakistan. Pakistan Journal of Botany. 2011; 43(5):2301-2310.
  22. Batish DR, Kaur M, Harminder P, Singh R, Kohli K. Phytotoxicity of a medicinal plant, *Anisomeles indica*, against *Phalaris minor* and its potential use as natural herbicide in wheat fields. Crop Protection. 2007; 26(7):948-952.
  23. Negi JS, Singh P, Joshi GP, Rawat MS, Bisht VK. Chemical constituents of *Asparagus*. Pharmacognosy Reviews. 2010; 4(8):215-220.
  24. Sajem AL, Gosai K. Traditional use of medicinal plants by the Jaintia tribes in North Cachar Hills district of Assam. Journal of Ethnobiology and Ethnomedicine. 2006; 2:2-7.
  25. Goyal RK, Singh J, Lal H. *Asparagus racemosus* - An update. Indian Journal of Medical Sciences. 2003; 57(9):408-414.
  26. Gautam M, Saha S, Bani S, Kaul A, Mishra S, Patil D *et al.* Immunomodulatory activity of *Asparagus racemosus* on systemic Th1/Th2 immunity: Implications for immunoadjuvant potential. Journal of Ethnopharmacology. 2009; 121(2):241-247.
  27. Mitra MP, Saumya D, Sanjita D, Kumar DM. Phytopharmacology of *Berberis aristata* DC: A review. Journal of Drug Delivery and Therapeutics. 2011; 1(2): 46-50.
  28. Uniyal SK, Singh KN, Jamwal P, Lal B. Traditional use of medicinal plants among the tribal communities of Chhota Bhangal, Western Himalaya. Journal of Ethnobiology and Ethnomedicine. 2006; 2(14):1-8.
  29. Singh M, Srivastava S, Rawat AK. Antimicrobial activities of Indian Berberis species. Fitoterapia. 2007; 78:574-576.
  30. Qureshi RA, Ghufran MA, Gilani SA, Yousaf Z, Abbas G, Batool A. Indigenous Medicinal Plants used by Local Women in Southern Himalayan Regions of Pakistan. Pakistan Journal of Botany. 2009; 41(1):19-25.
  31. Wang J, Dali K, Wang L, Lin-min L, Eric Jung-chi L. A survey of Chinese herbal ingredients with liver protection activities Chinese Medicine. 2007; 2(5):1-8.
  32. Adesina SK. Anticonvulsant Properties of the Roots of *Boerhaavia diffusa* Linnaeus. Quarterly journal of crude drug research. 2008; 17(2):84-86.
  33. Kumar PPNV, Pammi SVN, Kollu P, Satyanarayanan KV, Shameema U. Green synthesis and characterization of silver nanoparticles using *Boerhaavia diffusa* plant extract and their anti-bacterial activity. Industrial Crops and Products. 2014; 52:562-566.
  34. Patil HM, Bhaskar VV. Medicinal uses of plants by tribal medicine men of Nandurbar district in Maharashtra Explorer. Indian Journal of Natural Products and Resources. 2006; 5(2):125-130.
  35. Siddiqui MZ. *Boswellia Serrata*, A Potential Anti-inflammatory Agent: An Overview. Indian Journal of Pharmaceutical Sciences. 2011; 73(3):255-261.
  36. Borrelli F, Capasso F, Capasso R, Ascione V, Aviello G, Longo R *et al.* Effect of *Boswellia serrata* on intestinal motility in rodents, inhibition of diarrhoea without constipation. British journal of pharmacology. 2009; 148:553-560.
  37. Ekka NR, Dixit VK. Ethno-Pharmacognostical studies of Medicinal Plants of Jashpur District (Chhattisgarh). International Journal of Green Pharmacy. 2007; 1(1):2-4.
  38. Carmeliet P, Jain RK. Principles and mechanisms of vessel normalization for cancer and other angiogenic diseases. Nature Reviews Drug Discovery. 2011; 10:417-427.
  39. Choedon T, Shukla SK, Kumar V. Chemopreventive and anti-cancer properties of the aqueous extract of flowers of *Butea monosperma*. Journal of ethnopharmacology. 2010; 129(2):208-13.
  40. Pattanayak S, Nayak SS, Panda DP, Dinda SC, Shende V, Yadav A. Hepatoprotective Activity of Crude Flavonoids Extract of *Cajanus scarabaeoides* (L) in Paracetamol Intoxicated Albino Rats. Asian Journal of Pharmaceutical and Biological Research. 2011; 1(1):22-27.
  41. Rai AS, Rahman CH. Pharmacognostic standardization and phytochemical investigation of *Cajanus scarabaeoides* (L) Thouars. Research Journal of Pharmacognosy and Phytochemistry. 2018; 10(1):120-131.
  42. Sharma L, Agarwal G, Kumar A. Medicinal plants for skin and hair care. Indian Journal of Traditional Knowledge. 2003; 2(1):62-68.
  43. Hossain MS, Hanif A, Khan M, Bari S, Jahan R, Rahmatullah M. Ethnobotanical survey of the Tripura tribe of Bangladesh. American-Eurasian Journal of Sustainable Agriculture. 2009; 3(2):253-261.
  44. Kumar G, Karthik L, Kokati Rao KVB. *In vitro* anti-Candida activity of *Calotropis gigantean*. Journal of Pharmacy Research. 2010; 3(3):539-542.
  45. Parmar C, Kaushal MK. *Pyrus pashia*. In: Buch & Ham (Eds), Wild Fruits, Kalyani Publishers, New Delhi, India, 1982, 78-80.
  46. Singh S, Srivastava R, Choudhary S. Antifungal and HPLC analysis of the crude extracts of *Acorus calamus*, *Tinospora cordifolia* and *Celestrus paniculatus*. Journal of Agricultural Technology. 2010; 6(1):149-158.
  47. Dhiman AK. Conservation of some useful medicinal plants of Haridwar district in Uttaranchal state. In: Chopra AK, Khanna DR, Prasad G, Malik DS, Bhutiani R (ed.) Medicinal plants: Conservation, Cultivation and Utilization, 2007, 147-166.
  48. Shukla AN, Srivastava S, Rawat AKS. An ethnobotanical study of medicinal plants Rewa District Madhya Pradesh. Indian Journal of Traditional Knowledge. 2010; 9(1):191-202.



49. Mali RG. *Cleome viscosa* (wild mustard), A review on ethnobotany, phytochemistry, and pharmacology. *Journal of Pharmaceutical and Biological Sciences*. 2009; 48(1):105-112.
50. Ozturk M, Uysal I, Gucl S, Altundag E, Dogan Y, Baslar S. Medicinal Uses of Natural Dye-Yielding Plants in Turkey. *Research Journal of Textile and Apparel*. 2013; 17(2):69-80.
51. Silva BM, Andrade PB, Goncalves AC, Seabra RM, Oliveira MB. Influence of jam processing upon the contents of phenolics organic acids and free amino acids in quince fruit (*Cydonia oblonga* Miller). *European Food Research and Technology*. 2004; 218(4):385-389.
52. De Tommasi N, De Simone F, Pizza C, Mahmood N. Newtetracyclic sesterterpenes from *Cydonia vulgaris*. *Journal of Natural Products*. 1996; 59(3):267-270.
53. Oliveira AP, Pereira JA, Andrade PB, Valentão P, Seabra RM. Phenolic profile of *Cydonia oblonga* Miller leaves. *Journal of Agricultural and Food Chemistry*. 2007; 55(19):7926-7930.
54. Usmanghani K, Saeed A, Alam MT. *Indusyunic Medicine*. Uni of Karachi press, Karachi, 1997.
55. Duke JA, Bogenschutz-Godwin MJ, Ducealliar J, Duke PAK. *Handbook of Medicinal Herbs* (2nd edn). CRC Press, Boca Raton, 2002.
56. Prajapati ND, Purohit SS, Sharma AK, Kumar T. *A Handbook of Medicinal Plants*, Agrobios India, 2006, 184.
57. Bruin D, Baars E. *Citrus/Cydonia comp*, Use in general practice, A survey among anthroposophic physicians. LBI, Driebergen, The Netherlands, 2001.
58. Sakthi SS, Geetha M, Saranraj P. Pharmacological screening of *Datura metel* and *Acalypha indica* for its antifungal activity against pathogenic fungi. *International Journal of Pharmaceutical Sciences Review and Research*. 2011; 2(1):15-29.
59. Soni P, Siddiqui AA, Dwivedi J, Soni V. Pharmacological properties of *Datura stramonium* L. as a potential medicinal tree: An overview. *Asian Pacific Journal of Tropical Biomedicine*. 2012; 2(12):1002-1008.
60. Shah A, Marwat SK, Gohar F, Khan A, Bhatti KH, Amin M *et al*. Ethnobotanical Study of Medicinal Plants of Semi-Tribal Area of Makerwal & Gulla Khel (Lying between Khyber Pakhtunkhwa and Punjab Provinces). *American Journal of Plant Sciences*. 2013; 4:98-116.
61. Bozorgia M, Amin G, Shekarchi M, Rahimia R. Traditional medical uses of *Drimys* species in terms of phytochemistry, pharmacology and toxicology. *Journal of Traditional Chinese Medicine*. 2017; 37(1):124-139.
62. Pala NA, Negi AK, Todaria NP. Traditional uses of medicinal plants of Pauri Garhwal, Uttarakhand. *Nature and Science*. 2010; 8(6):57-61.
63. Rout SD, Panda T, Mishra N. Ethnomedicinal plants used to cure different diseases by tribal's of Mayurbhanj district of North Orissa. *Ethno-Med*. 2009; 3(1):27-32.
64. Mazumder PM, Farswan M, Parcha V. Hypoglycaemic effect of *Ficus arnottiana* Miq Bark extracts on streptozocin induced diabetes in rats. *Natural Product Radiance*. 2009; 8(5):478-482.
65. Ghosh R, Sharathchandra K, Rita S, Thokchom IS. Hypoglycaemic activity of *Ficus hispida* (bark) in normal and diabetic albino rats. *Indian Journal Pharmacology*. 2004; 36(4):222-225.
66. Aswar M, Aswar U, Watkar B, Vyas M, Wagh A. Anthelmintic activity of *Ficus bengalensis*. *International Journal of Green Pharmacy*. 2008; 2(3):170-172.
67. Ayinde BA, Omogbai EK, Amaechina FC. Pharmacognosy and hypotensive evaluation of *Ficus exasperate* Vahl (Moraceae) leaf. *Acta Poloniae Pharmaceutica*. 2007; 64(6):543-546.
68. Singh D, Singh B, Goel RK. Traditional uses, phytochemistry and pharmacology of *Ficus religiosa*: A review. *Journal of Ethnopharmacology*. 2011; 134(3):565-583.
69. Bhowmik D, Gopinath H, Kumar BP, Duraiavel S, Aravind G. Medicinal Uses of *Punica granatum* and its Health Benefits. *Journal of Pharmacognosy and Phytochemistry*. 2013; 1(5):2278-4136.
70. Mahmood A, Malik RN, Shinwari ZK, Mahmood A. Ethnobotanical survey of plants from Neelum, Azad Jammu and Kashmir, Pakistan. *Pakistan Journal of Botany*. 2011; 43:105-110.
71. Kanetkar P, Singhal R, Kamat M. *Gymnema sylvestre*: A Memoir. *Journal of Clinical Biochemistry and Nutrition*. 2007; 41(2):77-81.
72. Giri D, Tamta S, Pandey A. A review account on medicinal value of *Hedychium spicatum* Buch-Ham ex Sm: Vulnerable medicinal plant. *Journal of Medicinal Plants Research*. 2010; 4(25):2773-2777.
73. Venkatesh S, Reddy GD, Reddy BM. Antihyperglycemic Activity of *Helicteres isora* Roots in Alloxan-Induced Diabetic Rats. *Pharmaceutical Biology*. 2004; 41(5):347-350.
74. Chirayil CJ, Joy J, Mathew L, Mozetic M, Koetz J, Thomas S. Isolation and characterization of cellulose nanofibrils from *Helicteres isora* plant. *Industrial Crops and Products*. 2014; 59:27-34.
75. Gilani AH, Khan A, Khan A, Bashir S, Rahman N, Mandukhail SR. Pharmacological basis for the medicinal use of *Holarrhena antidysenterica* in gut motility disorders. *Journal of Pharmaceutical Biology*. 2010; 48(11):1240-1246.
76. Das RJ, Pathak K. Use of Indigenous Plants in Traditional Health Care Systems by Mishing Tribe of Dikhowmukh, Sivasagar District, Assam. *International Journal of Health and Medicine*. 2013; 1(3):50-57.
77. Shanmugasundaram P, Venkataraman S. Anti-Nociceptive Activity of *Hygrophila auriculata* (Schum) Heine. *African Journal of Traditional, Complementary and Alternative Medicines*. 2006; 2(1):62-69.
78. Hussain S, Ahmed N, Ansari Z. Preliminary studies on diuretic effect of *Hygrophila auriculata* (schum) heine in rats. *International Journal of Health Sciences Education*. 2009; 2(1):59-64.
79. Wang YC, Chuang YC, Hsu HW. The flavonoid, carotenoid and pectin content in peels of Citrus cultivated in Taiwan. *Food Chemistry*. 2008; 106:277-284.
80. Gordien YA, Gray IA, Franzblau GS, Seidel V. Antimycobacterial terpenoids from *Juniperus communis* L. (Cupressaceae). *Journal of Ethnopharmacology*. 2009; 126(3):500-505.
81. Miceli N, Trovato A, Dugo P, Cacciola F, Donato P, Marino A *et al*. Comparative Analysis of Flavonoid Profile, Antioxidant and Antimicrobial Activity of the Berries of *Juniperus communis* L var *communis* and *Juniperus communis* L var *saxatilis* Pall from Turkey. *Journal of Agricultural and Food Chemistry*. 2009; 57(15):6570-6577.
82. Rahmatullah M, Ferdousi D, Mollik MAH, Jahan R, Chowdhury MH, Haque WM. A survey of medicinal



- plants used by kavirajes of chalna area, Khulna district, Bangladesh. African Journal of Traditional, Complementary and Alternative Medicines. 2010; 7(2):91-97.
83. Biswas A, Haq WM, Akber M, Ferdausi D, Seraj S, Jahan FI, Chowdhury AR, Rahmatullah M. A survey of medicinal plants used by folk medicinal practitioners of Paschim Shawra and Palordi villages of Gaurnadi Upazila in Barisal district, Bangladesh. Journal of Sustainable Agriculture. 2011; 5:15-22.
  84. Al-Snafi AE. The pharmacology and medical importance of *Dolichos lablab* (*Lablab purpureus*) - A review. IOSR Journal of Pharmacy. 2017; 7(2):22-30.
  85. Rahmatullah M, Azam MNK, Rahman MM, Seraj S, Mahal MJ, Mou SM, Nasrin D, Khatun Z, Islam F, Chowdhury HM. A Survey of Medicinal Plants Used by Garo and Non-Garo Traditional Medicinal Practitioners in Two Villages of Tangail District, Bangladesh. American-Eurasian Journal of Sustainable Agriculture. 2011; 5(3):350-357.
  86. Anisuzzaman M, Rahman AHMM, Rashid M, Harun-Or-Rashid M, Naderuzzaman ATM, Islam AKMR. An Ethnobotanical Study of Madhupur, Tangail, Journal of Applied Sciences Research. 2007; 3(7):519-530.
  87. Rahman AHMM, Hossain MM, Islam AKMR. Taxonomy and Medicinal Uses of Angiosperm Weeds in the Wheat Field of Rajshahi, Bangladesh. Frontiers of Biological and Life Sciences. 2014; 2(1):8-11.
  88. Dhyani A. Exploring *Lilium polyphyllum* in Uttarakhand, India. The Lily Yearbook of North American Lily Society. Edt David Sims, 2007, 79-82.
  89. Ramana KV, Solomon AJR. Traditional and commercial uses of *Litsea glutinosa* (Lour) CB Robinson (Lauraceae). Journal of Medicinal Plants Research. 2017; 5(3):89-91.
  90. Sen A, Batra A. Evaluation of antimicrobial activity of different solvent extracts of medicinal plant: *Melia azedarach* L. International Journal of Current Pharmaceutical Research. 2012; 4(2):67-73.
  91. Naghibi F, Mosaddegh M, Mohammadi MM, Ghorbani A. Labiatae Family in folk Medicine in Iran. Journal of Ethnobiology and Ethnomedicine. 2005; 4(2):63-79.
  92. Tiwari GBGPK, Tadele K, Aramde F, Tiwari SC. Community Structure and Regeneration Potential of *Shorea robusta* Forest in Subtropical Submontane Zone of Garhwal Himalaya, India. Nature and Science. 2010; 8(1):68-74.
  93. Kumar A, Rana AC. Pharmagnotic and pharmacological profile of traditional medicinal plant: *Myrica nagi*. International Research Journal of Pharmacy. 2012; 3(12):32-37.
  94. Panthari P, Kharkwal H, Kharkwal H, Joshi DD. *Myrica nagi*: a review on active constituents, biological and therapeutic effects. International Journal of Pharmacy and Pharmaceutical Sciences. 2012; 4(5):38-42.
  95. Gusain YS, Khanduri VP. *Myrica esculenta* wild edible fruit of Indian Himalaya: need a sustainable approach for indigenous utilization. Ecology, Environment and Conservation. 2016; 22:267-270.
  96. Jeeva S, Lyndem FG, Sawian JT, Laloo RC, Mishra BP. *Myrica esculenta* Buch.- Ham. ex D Don - a potential ethnomedicinal species in a subtropical forest of Meghalaya, northeast India. Asian Pacific Journal of Tropical Biomedicine. 2011; 1(2):174-177.
  97. Nitha B, Meera CR, Janardhanan, KK. Anti-inflammatory and antitumour activities of cultured mycelium of morel mushroom, *Morchella esculenta*. Current Science. 2007; 92(2):235-239.
  98. Katkar KV, Suthar AC, Chauhan VS. The chemistry, pharmacologic, and therapeutic applications of *Polyalthia longifolia*. Pharmacognosy Reviews. 2010; 4(7):62-68.
  99. Deshmukh RD, Pokharkar RD, Takate SB, Gite VN. Amelioration of Ccl4-Induced Hepatosuppression by *Nyctanthes arbortristis* Linn Leaves in Wistar Albino Rats. Pharmacology online. 2007; 1:203-208.
  100. Khosravi AR, Shokri H, Kermani S, Dakhili M, Madani M, Parsa S. Antifungal properties of *Artemisia sieberi* and *Origanum vulgare* essential oils against *Candida glabrata* isolates obtained from patients with vulvovaginal candidiasis. Journal de Mycologie Médicale. 2011; 21(2):93-99.
  101. Lemhadri A, Zeggwagh NA, Maghrani M, Jouad H, Eddouks M. Anti-hyperglycaemic activity of the aqueous extract of *Origanum vulgare* growing wild in Tafilalet region. Journal of Ethnopharmacology. 2004; 92(2-3):251-256.
  102. Raghavendra MP, Satish S, Raveesha KA. Phytochemical analysis and antibacterial activity of *Oxalis corniculata*; a known medicinal plant. myScience. 2006; 1(1):72-78.
  103. Krishnaveni M, Mirunalin S. Amla-The Role of Ayurvedic Therapeutic Herb in Cancer. Asian Journal of Pharmaceutical and Clinical Research. 2011; 4(3):13-17.
  104. Bhide MM, Nitave SA. Roles of *Embllica officinalis* (amla) in medicine. World Journal of Pharmaceutical Sciences. 2014 3(6):604-615.
  105. Manoj P, Soniya EV, Banerjee NS, Ravichandran P. Recent studies on well-known spice, *Piper longum* Linn. Natural Product Radiance. 2004; 3(4):222-227.
  106. Samuelsen AB. The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review. Journal of Ethnopharmacology. 2000; 71(1-2):1-21.
  107. Jahan I, Mohammad SR, Mohammed ZR, Mohammad AK, Mohammad SI, Wahab A, Mohammad AR. Chemical and biological investigations of *Delonix regia* (Bojer ex Hook) Raf. Acta Pharmaceutica. 2010; 60:207-215.
  108. Tilak JC, Adhikari S, Devasagayam, TP. Antioxidant properties of *Plumbago zeylanica* an Indian medicinal plant and its active ingredient, plumbagin. Redox Report. 2004; 9(4):219-227.
  109. Wang YC, Huang TL. High-performance liquid chromatography for quantification of plumbagin, an anti-Helicobacter pylori compound of *Plumbago zeylanica* L. Journal of Chromatography A. 2005; 1094(1-2):99-104.
  110. Bhatt D, Kumar R, Tewari LM, Joshi GC. *Polygonatum cirrhifolium* Royle and *Polygonatum verticillatum* (L.) Allioni, Status assessment and medicinal uses in Uttarakhand, India. Journal of Medicinal Plants Research. 2014; 8(5):253-259.
  111. Khan H, Saeed M, Gilani AH, Khan MA, Dar A, Khan I. The antinociceptive activity of *Polygonatum verticillatum* rhizomes in pain models. Journal of Ethnopharmacology. 2010; 127:521-527.
  112. Sharma S, Satpathy G, Gupta RK. Nutritional, phytochemical, antioxidant and antimicrobial activity of *Prunus armenicus*. Journal of Pharmacognosy and Phytochemistry. 2014; 3(3):23-28.
  113. Yigit D, Yigit N, Mavi A. Antioxidant and antimicrobial activities of bitter and sweet apricot (*Prunus armeniaca*

- L) kernels. Brazilian Journal of Medical and Biological Research. 2009; 42(4):346-352.
114. Maria GM, Maria AN, Maria DA. Pomegranate (*Punica granatum* L) A medicinal plant with myriad biological properties - A short review. Journal of Medicinal Plants Research. 2010; 4(25):2836-2847.
115. Bisht VK, Kandari LS, Negi JS, Bhandari AK, Sundriyal RC. Traditional use of medicinal plants in district Chamoli, Uttarakhand, India. Journal of Medicinal Plants Research. 2013 7(15):918-929.
116. Saklani S, Chandra S. *In Vitro* Antimicrobial Activity, Nutritional Value, Antinutritional Value and Phytochemical Screening of *Pyracantha crenulata* Fruit. International Journal of Pharmaceutical Sciences Review and Research. 2014; 26(1):1-5.
117. Nielsen IL, Haren GR, Magnussen EL, Dragstead LO, Rasmussen SE. Quantification of anthocyanins in commercial black currant juices by simple high-performance liquid chromatography. Investigation of their pH stability and antioxidative potency. Journal of Agricultural and Food Chemistry. 2003; 51(20):5861-5866.
118. Olsson ME, Gustavsson KE, Andersson S, Nilsson A, Duan RD. Inhibition of cancer cell proliferation *in vitro* by fruit and berry extracts and correlations with antioxidant levels. Journal of Agricultural and Food Chemistry. 2004; 52(24):7264-7271.
119. Oyewole OI, Owoseni AA, Faboro EO. Studies on medicinal and toxicological properties of *Cajanus cajan*, *Ricinus communis* and *Thymus vulgaris* leaf extracts. Journal of Medicinal Plants Research. 2010; 4(19):2004–2006.
120. Ghrabi Z. A Guide to Medicinal Plants in North Africa: *Rosa canina* L. International Union for Conservation of Nature and Natural Resources: Malaga, Spain. 2005; 229-231.
121. Vântu S. *In vitro* multiplication of *Rosa canina* L. Analele științifice ale Universității Al I, Cuza Iași Tomul LVII, fasc 1, s II a., Biologie vegetala. 2011; 19-22.
122. Haq F, Ahmad H, Alam M. Traditional uses of medicinal plants of Nandiar Khuwarr catchment (District Battagram), Pakistan. Journal of Medicinal Plants Research. 2011; 5(1):39-48.
123. Ali ASE. The pharmacology and medical importance of *Dolichos lablab* (*Lablab purpureus*). Journal of Pharmaceutical Sciences. 2014; 7(2):22-30.
124. Munguia AR, Carrillo-Inungaray ML, Carranza-Álvarez C, Pimentel-González DJ, Alvarado-Sánchez B. Antioxidant activity, antimicrobial and effects in the immune system of plants and fruits extracts. Frontiers in Life Science. 2016; 9(2):90-98.
125. Prakash V, Aggarwal A. Traditional uses of ethnomedicinal plants of lower foothills of Himachal Pradesh-I. Indian journal of traditional knowledge. 2010; 9(3):519-521.
126. Tsewang TJ. Tibetan Medicinal Plants. Tibetan Medical Publications, India. 1995.
127. Ahmad S, Ullah F, Ayaz M, Sadiq A, Imran M. Antioxidant and anticholinesterase investigations of *Rumex hastatus* D. Don: potential effectiveness in oxidative stress and neurological disorders. Biological Research. 2015; 48(2):2-8.
128. Liang HX, Dai HQ, Fu HA, Dong XP, Adebayo AH, Zhang LX *et al.* Bioactive compounds from *Rumex* plants. Phytochemistry Letters. 2010; 3(4):181-184.
129. Kambhar SV. *Rumex vesicarius* L (Polygonaceae): An Overview. Journal of Global Ecology and Environment. 2014; 1(1):11-14.
130. Al Aboody MS. *In vitro* screening of Phytochemical, Antibacterial and Antioxidant activities of *Rumex vesicarius* L. International Journal of Current Microbiology and Applied Sciences. 2015; 4(9):884-893.
131. Mishra AP, Saklani S. *Satyrium nepalense*: A rare medicinal orchid of western Himalaya (India); phytochemical screening, antimicrobial evaluation and conservation studies. Indonesian Journal of Pharmacy. 2012; 23(3):162-170.
132. Silva RI, Melo GB, Melo VA, Antonioli AR, Michellone PR, Zucoloto S, Picinato MA, Franco CF. Effect of the aqueous extract of *Sida cordifolia* on liver regeneration after partial hepatectomy. Acta Cirúrgica Brasileira. 2006; 21(1):37-39.
133. Longo L, Vasapollo G. Extraction and Identification of Anthocyanins from *Smilax aspera* L. Berries. Food Chemistry. 2006; 94:226-231.
134. Saleem TSM, Chetty CM, Ramkanth S, Alagusundaram M, Gnanaprakash K. *Solanum nigrum* Linn- A Review. Pharmacognosy Reviews. 2009; 3(6):342-345.
135. Jain R, Sharma A, Gupta S, Sarethy IP, Gabrani R. *Solanum nigrum*: Current Perspectives on Therapeutic Properties. Alternative Medicine Review. 2011; 16(1):78-85.
136. Atanu FO, Ebiloma UG, Ajayi EI. A review of the pharmacological aspects of *Solanum nigrum* Linn, Biotechnology and Molecular Biology Reviews. 2011; 6(1):1-7.
137. Yousaf Z, Wang Y, Baydoun E. Phytochemistry and Pharmacological Studies on *Solanum torvum* Swartz. Journal of Applied Pharmaceutical Science. 2013; 3(4):152-160.
138. Nyeem MAB, Rashid AKMMU, Nowrose M, Hossain MA. *Solanum nigrum* (Maku): A review of pharmacological activities and clinical effects. International Journal of Applied Research. 2017; 3(1):12-17.
139. Rahmatullah M, Mollik MAH, Azam ATMA, Islam MR, Chowdhury MAM, Jahan R, Chowdhury MH, Rahman T. Ethnobotanical survey of the Santal tribe residing in Thakurgaon District, Bangladesh. American-Eurasian Journal of Sustainable Agriculture. 2009; 3:889–898.
140. Janbaz KH, Akhtar T, Saqib F, Imran I, Zia-Ul-Haq M, Jansakul C, De Feo V, Moga M. Pharmacological justification of use of *Solena heterophylla* Lour, in gastrointestinal, respiratory and vascular disorders. Journal of Translational Medicine. 2015; 13:2-8.
141. Joshi P, Dhawan V. *Swertia chirayita* – an overview. Current Science. 2005; 89(25):635-640.
142. Ayyanar M, Subash-Babu P. *Syzygium cumini* (L.) Skeels: A review of its phytochemical constituents and traditional uses. Asian Pacific Journal of Tropical Biomedicine. 2012; 2(3):240-246.
143. Bachaya HA, Iqbal Z, Khan MN, Jabbar A, Gilani AH, Din IU. *In vitro* and *in vivo* anthelmintic activity of *Terminalia arjuna* bark. International Journal of Agriculture and Biology. 2009; 11:273–278.
144. Gopinath K, Venkatesh KS, Ilangovan R, Sankaranarayanan K, Arumugam A. Green synthesis of gold nanoparticles from leaf extract of *Terminalia arjuna*, for the enhanced mitotic cell division and pollen germination activity. Industrial Crops and Products.

- 2013; 50:737-742.
145. Rahman MM, Mosadik MA, Wahed MI, Haque ME. Antimicrobial activity and cytotoxicity of *Trapa bispinosa*. *Fitoterapia*. 2000; 71:704-706.
  146. Hossan S, Agarwala B, Sarwar S, Karim M, Jahan R, Rahmatullah M). Traditional use of medicinal plants in Bangladesh to treat urinary tract infections and sexually transmitted diseases. *Ethnobotany Research and Applications*. 2010; 8:61-74.
  147. Shrestha P, Gautam R, Rana RB, Sthapit B. Home gardens and in situ conservation of plant genetic resources in farming systems. *Proceedings of the Second International Home Gardens Workshop, Witzenhausen, Federal Republic of Germany*. 2003; 105-108.
  148. Kalaivani T, Mathew L. Free radical scavenging activity from leaves of *Acacia nilotica* (L) Will ex Delile, an Indian medicinal tree. *Food and Chemical Toxicology*. 2010; 48:298-305.
  149. Shittu GA. *In vitro* antimicrobial and phytochemical activities of *Acacia nilotica* leaf extract. *Journal of Medicinal Plants Research*. 2010; 4(12):1232-1234.
  150. Agrawal S, Kulkarni GT, Sharma VN. A comparative study on the antioxidant activity of methanol extracts of acacia. *Advances in natural and applied sciences*. 2010; 4(1):78-84.
  151. Del WE. *In vitro* evaluation of peroxy radical scavenging capacity of water extract / fractions of *Acacia nilotica*. *African Journal of Biotechnology*. 2009; 8(7):1270-1272.
  152. Kaur K, Michael H, Arora S, Harkonen P, Kumar S. *In vitro* bioactivity-guided fractionation and characterization of polyphenolic inhibitory fractions from *Acacia nilotica* (L.) Willd ex Del. *Journal of Ethnopharmacology*. 2005; 99:353-630.
  153. Singh BN, Singh BR, Singh, RL, Prakash D, Sarma BK, Singh HB. Antioxidant and anti-quorum sensing activities of green pod of *Acacia nilotica* L. *Food and Chemical Toxicology*. 2009; 47:778-786.
  154. Singh R, Singh B, Singh S, Kumar N, Kumar S, Arora S. Antifree radical activities of kaempferol isolated from *Acacia nilotica* (L) Willd ex Del. *Toxicology in Vitro*. 2008; 22(8):1965-1970.
  155. Ali A, Akhtar N, Khan BA, Khan MS, Rasul A, Shahiq-UZ-Zaman, Khalid N, Waseem K, Mahmood T, Ali L. *Acacia nilotica*: A plant of multipurpose medicinal uses. *Journal of Medicinal Plants Research*. 2012; 6(9): 1492-1496.
  156. Turker AU, Campera ND. Biological activity of common mullein, a medicinal plant, *Journal of Ethnopharmacology*. 2002; 82(2-3):117-125.
  157. Shaheen S, Harun N, Khan F, Hussain RA, Ramzan S, Rani S, Khalid Z, Ahmad M, Zafar M. Comparative nutritional analysis between *Vigna radiata* and *Vigna mungo* of Pakistan. *African Journal of Biotechnology*. 2012; 11(25):6694-6702.
  158. Tandon VR. Medicinal uses and biological activities of *Vitex negundo*. *Indian Journal of Natural Products and Resources*. 2005; 4(3):162-165.
  159. Agarwal C, Singh RP, Agarwal R. Grape seed extract induces apoptotic death of human prostate carcinoma DU145 cells via caspases activation accompanied by dissipation of mitochondrial membrane potential and cytochrome c release. *Carcinogenesis*. 2002; 23(11):1869-1876.
  160. Bagchi D, Sen CK, Ray SD, Das DK, Bagchi M, Preuss HG, Vinson JA. Molecular mechanisms of cardioprotection by a novel grape seed proanthocyanidin extract. *Mutation Research*. 2003; 523-524:87-97.
  161. Bender DA, Bender AE. *A Dictionary of Food and Nutrition*, (4th edn), New York, Oxford University Press, USA 2005.
  162. Singh TP, Singh OM. Phytochemical and pharmacological profile of *Zanthoxylum armatum* DC - An overview. *Indian Journal of Natural Products and Resources*. 2011; 2(3):275-285.
  163. Zhang H, Jiang L, Ye S, Ye Y, Ren F. Systematic evaluation of antioxidant capacities of the ethalonic extract of different tissues of jujube (*Ziziphus jujube* Mill) from China. *Food and Chemical Toxicology*. 2010; 48(6):1461-1465.